

## Gaussian Quadrature with Legendre polynomials

- ▶ Legendre polynomials:  $P_0(x) = 1, P_1(x) = x,$

$$(n+1)P_{n+1}(x) = (2n+1)xP_n(x) - nP_{n-1}(x) \quad \text{for } n \geq 1.$$

## Gaussian Quadrature with Legendre polynomials

- ▶ Legendre polynomials:  $P_0(x) = 1, P_1(x) = x,$   
 $(n+1)P_{n+1}(x) = (2n+1)xP_n(x) - nP_{n-1}(x)$  for  $n \geq 1.$

- ▶ Gaussian quadrature

$$\int_{-1}^1 f(x) dx \approx c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n),$$

where  $x_1, x_2, \dots, x_n \in (-1, 1)$  are distinct roots of  $P_n(x).$

## Gaussian Quadrature with Legendre polynomials

- ▶ Legendre polynomials:  $P_0(x) = 1, P_1(x) = x,$   
 $(n+1)P_{n+1}(x) = (2n+1)xP_n(x) - nP_{n-1}(x)$  for  $n \geq 1.$

- ▶ Gaussian quadrature

$$\int_{-1}^1 f(x) dx \approx c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n),$$

where  $x_1, x_2, \dots, x_n \in (-1, 1)$  are distinct roots of  $P_n(x).$

- ▶ **DoP** =  $2n - 1$  with choice

$$c_i \stackrel{\text{def}}{=} \int_{-1}^1 L_i(x) dx = \int_{-1}^1 \left( \prod_{j \neq i} \frac{x - x_j}{x_i - x_j} \right) dx \quad \text{for } i = 1, \dots, n.$$

## Gaussian Quadrature with Legendre polynomials

- ▶ Legendre polynomials:  $P_0(x) = 1, P_1(x) = x,$   
 $(n+1)P_{n+1}(x) = (2n+1)xP_n(x) - nP_{n-1}(x)$  for  $n \geq 1.$

- ▶ Gaussian quadrature

$$\int_{-1}^1 f(x) dx \approx c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n),$$

where  $x_1, x_2, \dots, x_n \in (-1, 1)$  are distinct roots of  $P_n(x).$

- ▶ **DoP** =  $2n - 1$  with choice

$$c_i \stackrel{\text{def}}{=} \int_{-1}^1 L_i(x) dx = \int_{-1}^1 \left( \prod_{j \neq i} \frac{x - x_j}{x_i - x_j} \right) dx \quad \text{for } i = 1, \dots, n.$$

- ▶ Next: Estimate error in quadrature

## Hermite Interpolation, with Legendre polynomial $P_n(x)$

- Given Legendre roots  $x_1, x_2, \dots, x_n$  with

$$(x_1, f(x_1), f'(x_1)), (x_2, f(x_2), f'(x_2)), \dots, (x_n, f(x_n), f'(x_n)),$$

## Hermite Interpolation, with Legendre polynomial $P_n(x)$

- Given Legendre roots  $x_1, x_2, \dots, x_n$  with

$$(x_1, f(x_1), f'(x_1)), (x_2, f(x_2), f'(x_2)), \dots, (x_n, f(x_n), f'(x_n)),$$

- Interpolating polynomial  $H(x)$  of degree  $\leq 2n - 1$  satisfies

$$H(x_1) = f(x_1), \quad H'(x_1) = f'(x_1),$$

$$H(x_2) = f(x_2), \quad H'(x_2) = f'(x_2),$$

$$\vdots \quad \vdots$$

$$H(x_n) = f(x_n), \quad H'(x_n) = f'(x_n).$$

## Hermite Interpolation, with Legendre polynomial $P_n(x)$

- Given Legendre roots  $x_1, x_2, \dots, x_n$  with

$$(x_1, f(x_1), f'(x_1)), (x_2, f(x_2), f'(x_2)), \dots, (x_n, f(x_n), f'(x_n)),$$

- Interpolating polynomial  $H(x)$  of degree  $\leq 2n - 1$  satisfies

$$H(x_1) = f(x_1), \quad H'(x_1) = f'(x_1),$$

$$H(x_2) = f(x_2), \quad H'(x_2) = f'(x_2),$$

$$\vdots \quad \vdots$$

$$H(x_n) = f(x_n), \quad H'(x_n) = f'(x_n).$$

- Theorem:** For each  $x \in [a, b]$ , a number  $\xi(x)$  between  $x_1, x_2, \dots, x_n$  (hence  $\in (a, b)$ ) exists with

$$f(x) = H(x) + R(x), \quad R(x) = \frac{f^{(2n)}(\xi(x))}{(2n)!} (x-x_1)^2(x-x_2)^2 \cdots (x-x_n)^2.$$

## Hermite Interpolation, with Gaussian Quadrature

$$\int_{-1}^1 f(x) dx = \int_{-1}^1 H(x) dx + \int_{-1}^1 R(x) dx$$

## Hermite Interpolation, with Gaussian Quadrature

$$\int_{-1}^1 f(x) dx = \int_{-1}^1 H(x) dx + \int_{-1}^1 R(x) dx$$
$$\underbrace{\quad}_{\deg(H) \leq 2n-1} c_1 P(x_1) + c_2 P(x_2) + \cdots + c_n P(x_n) + \int_{-1}^1 R(x) dx$$

# Hermite Interpolation, with Gaussian Quadrature

$$\int_{-1}^1 f(x) dx = \int_{-1}^1 H(x) dx + \int_{-1}^1 R(x) dx$$
$$\underbrace{\quad}_{\deg(H) \leq 2n-1} c_1 P(x_1) + c_2 P(x_2) + \cdots + c_n P(x_n) + \int_{-1}^1 R(x) dx$$
$$\underbrace{\quad}_{\text{magic}} c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n)$$
$$+ \int_{-1}^1 \frac{f^{(2n)}(\xi(x))}{(2n)!} (x - x_1)^2 (x - x_2)^2 \cdots (x - x_n)^2 dx$$

# Hermite Interpolation, with Gaussian Quadrature

$$\int_{-1}^1 f(x) dx = \int_{-1}^1 H(x) dx + \int_{-1}^1 R(x) dx$$
$$\underbrace{\quad}_{\deg(H) \leq 2n-1} c_1 P(x_1) + c_2 P(x_2) + \cdots + c_n P(x_n) + \int_{-1}^1 R(x) dx$$
$$\underbrace{\quad}_{\text{magic}} c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n)$$
$$+ \int_{-1}^1 \frac{f^{(2n)}(\xi(x))}{(2n)!} (x - x_1)^2 (x - x_2)^2 \cdots (x - x_n)^2 dx$$
$$= c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n)$$
$$+ \frac{f^{(2n)}(\xi)}{(2n)!} \int_{-1}^1 (x - x_1)^2 (x - x_2)^2 \cdots (x - x_n)^2 dx$$

# Hermite Interpolation, with Gaussian Quadrature

$$\int_{-1}^1 f(x) dx = \int_{-1}^1 H(x) dx + \int_{-1}^1 R(x) dx$$
$$\stackrel{\deg(H) \leq 2n-1}{=} c_1 P(x_1) + c_2 P(x_2) + \cdots + c_n P(x_n) + \int_{-1}^1 R(x) dx$$
$$\stackrel{\text{magic}}{=} c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n)$$
$$+ \int_{-1}^1 \frac{f^{(2n)}(\xi(x))}{(2n)!} (x - x_1)^2 (x - x_2)^2 \cdots (x - x_n)^2 dx$$
$$= c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n)$$
$$+ \frac{f^{(2n)}(\xi)}{(2n)!} \int_{-1}^1 (x - x_1)^2 (x - x_2)^2 \cdots (x - x_n)^2 dx$$
$$\stackrel{\text{def}}{=} c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n) + \mathbf{R}$$

# Gaussian Quadrature Error Estimate

$$R = \frac{f^{(2n)}(\xi)}{(2n)!} \int_{-1}^1 (x - x_1)^2(x - x_2)^2 \cdots (x - x_n)^2 dx$$

## Gaussian Quadrature Error Estimate

$$\begin{aligned}\mathbf{R} &= \frac{f^{(2n)}(\xi)}{(2n)!} \int_{-1}^1 (x - x_1)^2(x - x_2)^2 \cdots (x - x_n)^2 dx \\ &= \frac{2^{2n} (n!)^3 (n-1)!}{(2n+1)! (2n)! (2n-1)!} f^{(2n)}(\xi) = O\left(\frac{4^{-n} |f^{(2n)}(\xi)|}{(2n)!}\right).\end{aligned}$$

# Gaussian Quadrature Error Estimate

$$\begin{aligned} R &= \frac{f^{(2n)}(\xi)}{(2n)!} \int_{-1}^1 (x - x_1)^2(x - x_2)^2 \cdots (x - x_n)^2 dx \\ &= \frac{2^{2n} (n!)^3 (n-1)!}{(2n+1)! (2n)! (2n-1)!} f^{(2n)}(\xi) = O\left(\frac{4^{-n} |f^{(2n)}(\xi)|}{(2n)!}\right). \end{aligned}$$


$$\int_{-1}^1 f(x) dx = c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n) + O\left(\frac{4^{-n} |f^{(2n)}(\xi)|}{(2n)!}\right).$$

Rapid convergence for smooth functions

## Composite Simpson's Rule

$$(n = 2m, x_j = a + j h, h = \frac{b-a}{n}, 0 \leq j \leq n)$$

$$\int_a^b f(x) dx = \frac{h}{3} \left( f(a) + 2 \sum_{j=1}^{m-1} f(x_{2j}) + 4 \sum_{j=1}^m f(x_{2j-1}) + f(b) \right) - \frac{(b-a)h^4}{180} f^{(4)}(\mu)$$



# Limitations of Gaussian Quadrature

Simpson/Trapezoidal:


- ▶ Composite rules:
  - ▶ Adding more EQUI-SPACED points.
- ▶ Romberg extrapolation:
  - ▶ Obtaining higher order rules from lower order rules.
- ▶ Adaptive quadratures:
  - ▶ Adding more points ONLY WHEN NECESSARY.

Gaussian Quadrature:

- ▶ points different for different  $n$ .

Gaussian Quadrature good for given  $n$ ,  
not as good for given tolerance.

Double Integral  $\int \int_R f(x, y) dA$



$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$$

## Double Integral = Integral of Integral

$$\begin{aligned}\int \int_R f(x, y) dA &= \int_a^b \left( \int_c^d f(x, y) dy \right) dx \\ &= \int_a^b g(x) dx,\end{aligned}$$

where  $g(x) \stackrel{\text{def}}{=} \int_c^d f(x, y) dy.$

### Approach

- ▶ Approximate  $\int_a^b g(x) dx$  with quadrature.
- ▶ For any given  $x_i$ , approximate  $g(x_i)$  with quadrature.

$$\int \int_R f(x, y) dA = \int_a^b g(x) dx, \quad g(x) = \int_c^d f(x, y) dy$$

► Approximate  $\int_a^b g(x) dx$  with  $n$ -point quadrature:

$$\int_a^b g(x) dx = c_1 g(x_1) + c_2 g(x_2) + \cdots + c_n g(x_n) + \mathbf{R}(g)$$

$$\int \int_R f(x, y) dA = \int_a^b g(x) dx, \quad g(x) = \int_c^d f(x, y) dy$$

- ▶ Approximate  $\int_a^b g(x) dx$  with  $n$ -point quadrature:

$$\int_a^b g(x) dx = c_1 g(x_1) + c_2 g(x_2) + \cdots + c_n g(x_n) + \mathbf{R}(g)$$

- ▶ For  $1 \leq i \leq n$ , approximate  $g(x_i)$  with  $m$ -point quadrature:

$$\int_c^d f(x_i, y) dy = \widehat{c}_1 f(x_i, y_1) + \widehat{c}_2 f(x_i, y_2) + \cdots + \widehat{c}_m f(x_i, y_m) + \widehat{\mathbf{R}}(f(x_i, \cdot)).$$

$$\int \int_R f(x, y) dA = \int_a^b g(x) dx, \quad g(x) = \int_c^d f(x, y) dy$$

- ▶ Approximate  $\int_a^b g(x) dx$  with  $n$ -point quadrature:

$$\int_a^b g(x) dx = c_1 g(x_1) + c_2 g(x_2) + \cdots + c_n g(x_n) + \mathbf{R}(g)$$

- ▶ For  $1 \leq i \leq n$ , approximate  $g(x_i)$  with  $m$ -point quadrature:

$$\int_c^d f(x_i, y) dy = \widehat{c}_1 f(x_i, y_1) + \widehat{c}_2 f(x_i, y_2) + \cdots + \widehat{c}_m f(x_i, y_m) + \widehat{\mathbf{R}}(f(x_i, \cdot)).$$

$$\begin{aligned} \int \int_R f(x, y) dA &= \left( \sum_{i=1}^n c_i g(x_i) \right) + \mathbf{R}(g) \\ &= \left( \sum_{i=1}^n c_i \left( \left( \sum_{j=1}^m \widehat{c}_j f(x_i, y_j) \right) + \widehat{\mathbf{R}}(f(x_i, \cdot)) \right) \right) + \mathbf{R}(g) \end{aligned}$$

$$\int \int_R f(x, y) dA = \int_a^b g(x) dx, \quad g(x) = \int_c^d f(x, y) dy$$

► Approximate  $\int_a^b g(x) dx$  with  $n$ -point quadrature:

$$\int_a^b g(x) dx = c_1 g(x_1) + c_2 g(x_2) + \cdots + c_n g(x_n) + \mathbf{R}(g)$$

► For  $1 \leq i \leq n$ , approximate  $g(x_i)$  with  $m$ -point quadrature:

$$\int_c^d f(x_i, y) dy = \widehat{c}_1 f(x_i, y_1) + \widehat{c}_2 f(x_i, y_2) + \cdots + \widehat{c}_m f(x_i, y_m) + \widehat{\mathbf{R}}(f(x_i, \cdot)).$$

$$\begin{aligned} \int \int_R f(x, y) dA &= \left( \sum_{i=1}^n c_i g(x_i) \right) + \mathbf{R}(g) \\ &= \left( \sum_{i=1}^n c_i \left( \left( \sum_{j=1}^m \widehat{c}_j f(x_i, y_j) \right) + \widehat{\mathbf{R}}(f(x_i, \cdot)) \right) \right) + \mathbf{R}(g) \\ &= \left( \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) \right) + \left( \sum_{i=1}^n c_i \widehat{\mathbf{R}}(f(x_i, \cdot)) \right) + \mathbf{R}(g). \end{aligned}$$

$$\int \int_R f(x, y) dA = \int_a^b g(x) dx, \quad g(x) = \int_c^d f(x, y) dy$$

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d.\}$$

$$\begin{aligned}\int_a^b g(x) dx &= c_1 g(x_1) + c_2 g(x_2) + \cdots + c_n g(x_n) + \mathbf{R}(g) \\ \int_c^d f(x_i, y) dy &= \widehat{c}_1 f(x_1, y_1) + \widehat{c}_2 f(x_i, y_2) + \cdots + \widehat{c}_m f(x_i, y_m) \\ &\quad + \widehat{\mathbf{R}}(f(x_i, \cdot))\end{aligned}$$

$$\int \int_R f(x, y) dA = \int_a^b g(x) dx, \quad g(x) = \int_c^d f(x, y) dy$$

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d.\}$$

$$\int_a^b g(x) dx = c_1 g(x_1) + c_2 g(x_2) + \cdots + c_n g(x_n) + \mathbf{R}(g)$$

$$\begin{aligned} \int_c^d f(x_i, y) dy &= \widehat{c}_1 f(x_1, y_1) + \widehat{c}_2 f(x_i, y_2) + \cdots + \widehat{c}_m f(x_i, y_m) \\ &\quad + \widehat{\mathbf{R}}(f(x_i, \cdot)) \end{aligned}$$

$$\begin{aligned} \int \int_R f(x, y) dA &= \left( \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) \right) + \left( \sum_{i=1}^n c_i \widehat{\mathbf{R}}(f(x_i, \cdot)) \right) + \mathbf{R}(g) \\ &\approx \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j). \end{aligned}$$

- ▶ Double integral quadrature is a double sum.
- ▶ Need to work out total error for any given quadrature.

$$\int \int_R f(x, y) dA = \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c_j} f(x_i, y_j) + \sum_{i=1}^n c_i \widehat{\mathbf{R}}(f(x_i, \cdot)) + \mathbf{R}(g)$$

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d.\}$$

**Example**, Simpson's Rule with  $m = n = 3$ :


- ▶ Simpson's Rule on  $[a, b]$ :  $(x_1, x_2, x_3) = (a, \frac{a+b}{2}, b)$ .
- ▶ Simpson's Rule on  $[c, d]$ :  $(y_1, y_2, y_3) = (c, \frac{c+d}{2}, d)$ .

$$\int \int_R f(x, y) dA = \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c_j} f(x_i, y_j) + \sum_{i=1}^n c_i \widehat{\mathbf{R}}(f(x_i, \cdot)) + \mathbf{R}(g)$$

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$$

**Example**, Simpson's Rule with  $m = n = 3$ :

- ▶ Simpson's Rule on  $[a, b]$ :  $(x_1, x_2, x_3) = (a, \frac{a+b}{2}, b)$ .
- ▶ Simpson's Rule on  $[c, d]$ :  $(y_1, y_2, y_3) = (c, \frac{c+d}{2}, d)$ .



$$\int \int_R f(x, y) dA = \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) + \sum_{i=1}^n c_i \widehat{\mathbf{R}}(f(x_i, \cdot)) + \mathbf{R}(g)$$

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}.$$

**Example,**  $m = n = 3$ :

- ▶ Simpson's Rule on  $[a, b]$ :  $(x_1, x_2, x_3) = (a, \frac{a+b}{2}, b)$ .

$$\int_a^b g(x) dx = c_1 g(x_1) + c_2 g(x_2) + c_3 g(x_3) - \frac{h^5}{90} g^{(4)}(\xi),$$

$$h = \frac{b-a}{2}, \quad (c_1, c_2, c_3) = \frac{h}{3} (1, 4, 1).$$

- ▶ Simpson's Rule on  $[c, d]$ :  $(y_1, y_2, y_3) = (c, \frac{c+d}{2}, d)$ .

$$\int_c^d f(x_i, y) dy = \widehat{c}_1 f(x_i, y_1) + \widehat{c}_2 f(x_i, y_2) + \widehat{c}_3 f(x_i, y_3) - \frac{k^5}{90} \frac{\partial^4 f}{\partial^4 y}(x_i, \eta_i),$$

$$k = \frac{d-c}{2}, \quad (\widehat{c}_1, \widehat{c}_2, \widehat{c}_3) = \frac{k}{3} (1, 4, 1).$$

$$\begin{aligned}
\int \int_R f(x, y) dA &= \left( \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) \right) - \frac{k^5}{90} \left( \sum_{i=1}^m c_i \frac{\partial^4 f}{\partial^4 y}(x_i, \eta_i) \right) \\
&\quad - \frac{h^5}{90} \int_a^b \frac{\partial^4 f}{\partial^4 x}(\xi, y) dy
\end{aligned}$$

$$\begin{aligned}
\int \int_R f(x, y) dA &= \left( \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) \right) - \frac{k^5}{90} \left( \sum_{i=1}^m c_i \frac{\partial^4 f}{\partial^4 y}(x_i, \eta_i) \right) \\
&\quad - \frac{h^5}{90} \int_a^b \frac{\partial^4 f}{\partial^4 x}(\xi, y) dy \\
&= \left( \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) \right) - \frac{k^5}{90} \left( \sum_{i=1}^m c_i \right) \frac{\partial^4 f}{\partial^4 y}(\widehat{\xi}, \widehat{\eta}) \\
&\quad - \frac{h^5}{90} (b - a) \frac{\partial^4 f}{\partial^4 x}(\xi, \eta)
\end{aligned}$$

$$\begin{aligned}
\int \int_R f(x, y) dA &= \left( \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) \right) - \frac{k^5}{90} \left( \sum_{i=1}^m c_i \frac{\partial^4 f}{\partial^4 y}(x_i, \eta_i) \right) \\
&\quad - \frac{h^5}{90} \int_a^b \frac{\partial^4 f}{\partial^4 x}(\xi, y) dy \\
&= \left( \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) \right) - \frac{k^5}{90} \left( \sum_{i=1}^m c_i \right) \frac{\partial^4 f}{\partial^4 y}(\widehat{\xi}, \widehat{\eta}) \\
&\quad - \frac{h^5}{90} (b-a) \frac{\partial^4 f}{\partial^4 x}(\xi, \eta) \\
&= \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) \\
&\quad - \frac{(b-a)(d-c)}{180} \left( k^4 \frac{\partial^4 f}{\partial^4 y}(\widehat{\xi}, \widehat{\eta}) + h^4 \frac{\partial^4 f}{\partial^4 x}(\xi, \eta) \right).
\end{aligned}$$

$$\int \int_R f(x, y) \, dA = \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) + \sum_{i=1}^n c_i \widehat{\mathbf{R}}(f(x_i, \cdot)) + \mathbf{R}(g)$$

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d.\}$$

**Example:** Composite Simpson Rules.

$$\int \int_R f(x, y) dA = \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) + \sum_{i=1}^n c_i \widehat{\mathbf{R}}(f(x_i, \cdot)) + \mathbf{R}(g)$$

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$$

**Example:** Composite Simpson Rules.

- ▶ Composite Simpson on  $[a, b]$ ,  $x_i = a + (i-1)h$ ,  $1 \leq i \leq n$ ,  $h = \frac{b-a}{n-1}$ .

$$\mathbf{R}(g) = -\frac{(b-a)h^4}{180} \int_c^d \frac{\partial^4 f}{\partial^4 x}(\xi, y) dy = -\frac{(b-a)(d-c)h^4}{180} \frac{\partial^4 f}{\partial^4 x}(\xi, \eta).$$

$$\int \int_R f(x, y) dA = \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j) + \sum_{i=1}^n c_i \widehat{\mathbf{R}}(f(x_i, \cdot)) + \mathbf{R}(g)$$

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$$

**Example:** Composite Simpson Rules.

- ▶ Composite Simpson on  $[a, b]$ ,  $x_i = a + (i-1)h$ ,  $1 \leq i \leq n$ ,  $h = \frac{b-a}{n-1}$ .

$$\mathbf{R}(g) = -\frac{(b-a)h^4}{180} \int_c^d \frac{\partial^4 f}{\partial^4 x}(\xi, y) dy = -\frac{(b-a)(d-c)h^4}{180} \frac{\partial^4 f}{\partial^4 x}(\xi, \eta).$$

- ▶ Composite Simpson on  $[c, d]$ ,  $y_j = c + (j-1)k$ ,  $1 \leq j \leq m$ ,  $k = \frac{d-c}{m-1}$ .

$$\begin{aligned}\widehat{\mathbf{R}}(f(x_i, \cdot)) &= -\frac{(d-c)k^4}{180} \frac{\partial^4 f}{\partial^4 y}(x_i, \eta_i) \\ \sum_{i=1}^n c_i \widehat{\mathbf{R}}(f(x_i, \cdot)) &= -\frac{(d-c)k^4}{180} \left( \sum_{i=1}^n c_i \right) \frac{\partial^4 f}{\partial^4 y}(\widehat{\xi}, \widehat{\eta}) \\ &= -\frac{(b-a)(d-c)k^4}{180} \frac{\partial^4 f}{\partial^4 y}(\widehat{\xi}, \widehat{\eta}).\end{aligned}$$

## Error Estimate, Double Integral with Composite Simpson

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$$


$$\begin{aligned} \int \int_R f(x, y) dA &= \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \\ &\quad - \frac{(b-a)(d-c)}{180} \left( k^4 \frac{\partial^4 f}{\partial^4 y}(\hat{\xi}, \hat{\eta}) + h^4 \frac{\partial^4 f}{\partial^4 x}(\xi, \eta) \right), \end{aligned}$$

$$h = \frac{b-a}{n-1}, \quad k = \frac{d-c}{m-1}.$$

Example:  $\int \int_R \log(x + 2y) dA$  with  $n = 7, m = 5$

$$R = \{(x, y) \mid 1.2 \leq x \leq 2.4, \quad 0.2 \leq y \leq 1.\}$$

Simpson Rule for double integral,  $n = 7, m = 5$



Example:  $\int \int_R \log(x + 2y) dA$  with  $n = 7, m = 5$

$$R = \{(x, y) \mid 1.2 \leq x \leq 2.4, \quad 0.2 \leq y \leq 1, \}$$

Example:  $\int \int_R \log(x + 2y) dA$  with  $n = 7, m = 5$

$$R = \{(x, y) \mid 1.2 \leq x \leq 2.4, \quad 0.2 \leq y \leq 1, \}$$

$$h = \frac{2.4 - 1.2}{7 - 1} = 0.2, \quad k = \frac{1 - 0.2}{5 - 1} = 0.2.$$

Example:  $\int \int_R \log(x + 2y) dA$  with  $n = 7, m = 5$

$$R = \{(x, y) \mid 1.2 \leq x \leq 2.4, \quad 0.2 \leq y \leq 1, \}$$

$$h = \frac{2.4 - 1.2}{7 - 1} = 0.2, \quad k = \frac{1 - 0.2}{5 - 1} = 0.2.$$

$$\frac{\partial^4 f}{\partial^4 x} = -\frac{6}{(x + 2y)^4}, \quad \frac{\partial^4 f}{\partial^4 y} = -\frac{96}{(x + 2y)^4}.$$

$$\left| \frac{\partial^4 f}{\partial^4 x} \right| \leq \frac{6}{(1.2 + 2 \times 0.2)^4} \approx 0.91553 \quad \text{for } (x, y) \in R,$$

$$\left| \frac{\partial^4 f}{\partial^4 y} \right| \leq \frac{96}{(1.2 + 2 \times 0.2)^4} \approx 14.648.$$

$$\begin{aligned}\text{Quad_Error} &= \frac{(b-a)(d-c)}{180} \left| k^4 \frac{\partial^4 f}{\partial^4 y}(\hat{\xi}, \hat{\eta}) + h^4 \frac{\partial^4 f}{\partial^4 x}(\xi, \eta) \right| \\ &\leq \frac{(2.4 - 1.2)(1 - 0.2)}{180} (0.2^4 \times 14.648 + 0.2^4 \times 0.91553)\end{aligned}$$

Example:  $\int \int_R \log(x + 2y) dA$  with  $n = 7, m = 5$

$$R = \{(x, y) \mid 1.2 \leq x \leq 2.4, \quad 0.2 \leq y \leq 1, \}$$

$$h = \frac{2.4 - 1.2}{7 - 1} = 0.2, \quad k = \frac{1 - 0.2}{5 - 1} = 0.2.$$

$$\frac{\partial^4 f}{\partial^4 x} = -\frac{6}{(x + 2y)^4}, \quad \frac{\partial^4 f}{\partial^4 y} = -\frac{96}{(x + 2y)^4}.$$

$$\left| \frac{\partial^4 f}{\partial^4 x} \right| \leq \frac{6}{(1.2 + 2 \times 0.2)^4} \approx 0.91553 \quad \text{for } (x, y) \in R,$$

$$\left| \frac{\partial^4 f}{\partial^4 y} \right| \leq \frac{96}{(1.2 + 2 \times 0.2)^4} \approx 14.648.$$

$$\begin{aligned}\text{Quad_Error} &= \frac{(b-a)(d-c)}{180} \left| k^4 \frac{\partial^4 f}{\partial^4 y}(\hat{\xi}, \hat{\eta}) + h^4 \frac{\partial^4 f}{\partial^4 x}(\xi, \eta) \right| \\ &\leq \frac{(2.4 - 1.2)(1 - 0.2)}{180} (0.2^4 \times 14.648 + 0.2^4 \times 0.91553) \\ &\approx 1.328 \times 10^{-4}.\end{aligned}$$

Example:  $\int \int_R \log(x + 2y) dA$  with  $n = 7, m = 5$

$$\int \int_R \log(x + 2y) dA = 1.0360481\cdots$$

Example:  $\int \int_R \log(x + 2y) dA$  with  $n = 7, m = 5$

$$\int \int_R \log(x + 2y) dA = 1.0360481\cdots$$

$$\begin{aligned}\int \int_R \log(x + 2y) dA &\approx \sum_{i=1}^7 \sum_{j=1}^5 c_i \hat{c}_j f(x_i, y_j) \\ &= 1.0360327\cdots.\end{aligned}$$

Example:  $\int \int_R \log(x + 2y) dA$  with  $n = 7, m = 5$

$$\int \int_R \log(x + 2y) dA = 1.0360481\cdots$$

$$\begin{aligned}\int \int_R \log(x + 2y) dA &\approx \sum_{i=1}^7 \sum_{j=1}^5 c_i \hat{c}_j f(x_i, y_j) \\ &= 1.0360327\cdots.\end{aligned}$$

Therefore

$$\left| \int \int_R \log(x + 2y) dA - \left( \sum_{i=1}^7 \sum_{j=1}^5 c_i \hat{c}_j f(x_i, y_j) \right) \right| = 1.546 \times 10^{-5}$$

Example:  $\int \int_R \log(x + 2y) dA$  with  $n = 7, m = 5$

$$\int \int_R \log(x + 2y) dA = 1.0360481\cdots$$

$$\begin{aligned}\int \int_R \log(x + 2y) dA &\approx \sum_{i=1}^7 \sum_{j=1}^5 c_i \hat{c}_j f(x_i, y_j) \\ &= 1.0360327\cdots.\end{aligned}$$

Therefore

$$\begin{aligned}\left| \int \int_R \log(x + 2y) dA - \left( \sum_{i=1}^7 \sum_{j=1}^5 c_i \hat{c}_j f(x_i, y_j) \right) \right| &= 1.546 \times 10^{-5} \\ &< 1.328 \times 10^{-4}. \\ &\quad (\text{Error Estimate})\end{aligned}$$

## 2 Dimensional Gaussian Quadratures

$$\begin{aligned} R &= \{(x, y) \mid a \leq x \leq b, c \leq y \leq d.\} \\ \int \int_R f(x, y) dA &= \int_a^b \left( \int_c^d f(x, y) dy \right) dx. \end{aligned}$$

## 2 Dimensional Gaussian Quadratures

$$R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$$
$$\int \int_R f(x, y) dA = \int_a^b \left( \int_c^d f(x, y) dy \right) dx.$$

Perform change of variables

$$x = \frac{a+b}{2} + \frac{b-a}{2} u, \quad y = \frac{c+d}{2} + \frac{d-c}{2} v \quad \text{for } u, v \in [-1, 1].$$

Double integral becomes

$$\int \int_R f(x, y) dA = \frac{(b-a)(d-c)}{4} \int_{-1}^1 \hat{g}(u) du, \quad \text{where}$$

$$\hat{g}(u) \stackrel{\text{def}}{=} \int_{-1}^1 f \left( \frac{a+b}{2} + \frac{b-a}{2} u, \frac{c+d}{2} + \frac{d-c}{2} v \right) dv.$$

$$\int \int_R f(x, y) dA = \frac{(b-a)(d-c)}{4} \int_{-1}^1 \widehat{g}(u) du$$

$$\widehat{g}(u) = \int_{-1}^1 f\left(\frac{a+b}{2} + \frac{b-a}{2} u, \frac{c+d}{2} + \frac{d-c}{2} v\right) dv.$$

- $n$ -point Gaussian quadrature for  $\int_{-1}^1 \widehat{g}(u) du$ :

$$\int_{-1}^1 \widehat{g}(u) du \approx c_1 \widehat{g}(u_1) + c_2 \widehat{g}(u_2) + \cdots + c_n \widehat{g}(u_n).$$

$$\int \int_R f(x, y) dA = \frac{(b-a)(d-c)}{4} \int_{-1}^1 \widehat{g}(u) du$$

$$\widehat{g}(u) = \int_{-1}^1 f\left(\frac{a+b}{2} + \frac{b-a}{2} u, \frac{c+d}{2} + \frac{d-c}{2} v\right) dv.$$

- $n$ -point Gaussian quadrature for  $\int_{-1}^1 \widehat{g}(u) du$ :

$$\int_{-1}^1 \widehat{g}(u) du \approx c_1 \widehat{g}(u_1) + c_2 \widehat{g}(u_2) + \cdots + c_n \widehat{g}(u_n).$$

- For  $1 \leq i \leq n$ , let

$$x_i = \frac{a+b}{2} + \frac{b-a}{2} u_i, \quad \widehat{g}(u_i) = \int_{-1}^1 f\left(x_i, \frac{c+d}{2} + \frac{d-c}{2} v\right) dv.$$

$$\int \int_R f(x, y) dA = \frac{(b-a)(d-c)}{4} \int_{-1}^1 \widehat{g}(u) du$$

$$\widehat{g}(u) = \int_{-1}^1 f\left(\frac{a+b}{2} + \frac{b-a}{2} u, \frac{c+d}{2} + \frac{d-c}{2} v\right) dv.$$

- $n$ -point Gaussian quadrature for  $\int_{-1}^1 \widehat{g}(u) du$ :

$$\int_{-1}^1 \widehat{g}(u) du \approx c_1 \widehat{g}(u_1) + c_2 \widehat{g}(u_2) + \cdots + c_n \widehat{g}(u_n).$$

- For  $1 \leq i \leq n$ , let

$$x_i = \frac{a+b}{2} + \frac{b-a}{2} u_i, \quad \widehat{g}(u_i) = \int_{-1}^1 f\left(x_i, \frac{c+d}{2} + \frac{d-c}{2} v\right) dv.$$

- $m$ -point Gaussian quadrature for  $\widehat{g}(u_i)$ ,

$$\begin{aligned} \widehat{g}(u_i) &\approx \widehat{c}_1 f\left(x_i, \frac{c+d}{2} + \frac{d-c}{2} v_1\right) + \cdots + \widehat{c}_m f\left(x_i, \frac{c+d}{2} + \frac{d-c}{2} v_m\right) \\ &= \widehat{c}_1 f(x_i, y_1) + \cdots + \widehat{c}_m f(x_i, y_m), \quad y_j \stackrel{\text{def}}{=} \frac{c+d}{2} + \frac{d-c}{2} v_j. \end{aligned}$$

$$\int \int_R f(x, y) dA = \frac{(b-a)(d-c)}{4} \int_{-1}^1 \widehat{g}(u) du$$

$$\begin{aligned}\int_{-1}^1 \widehat{g}(u) du &\approx c_1 \widehat{g}(u_1) + c_2 \widehat{g}(u_2) + \cdots + c_n \widehat{g}(u_n) \\ \widehat{g}(u_i) &\approx \widehat{c}_1 f(x_i, y_1) + \cdots + \widehat{c}_m f(x_i, y_m).\end{aligned}$$

So we have Gaussian quadrature for double integral:

$$\int \int_R f(x, y) dA \approx \frac{(b-a)(d-c)}{4} \sum_{i=1}^n \sum_{j=1}^m c_i \widehat{c}_j f(x_i, y_j).$$

Example:  $\int \int_R \log(x + 2y) dA = 1.036 \dots$ ,  $n = 7, m = 5$

- ▶ Gaussian quadrature approximation

$$\int \int_R f(x, y) dA \approx \frac{(b-a)(d-c)}{4} \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \approx 1.03604817065 \dots$$

$$\left| \int \int_R f(x, y) dA - \frac{(b-a)(d-c)}{4} \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \right| \approx 6.4 \times 10^{-10}.$$

Example:  $\int \int_R \log(x + 2y) dA = 1.036 \dots$ ,  $n = 7, m = 5$

- ▶ Gaussian quadrature approximation

$$\int \int_R f(x, y) dA \approx \frac{(b-a)(d-c)}{4} \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \approx 1.03604817065 \dots$$

$$\left| \int \int_R f(x, y) dA - \frac{(b-a)(d-c)}{4} \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \right| \approx 6.4 \times 10^{-10}.$$

- ▶ Simpson rule approximation

$$\int \int_R f(x, y) dA \approx \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \approx 1.03603270963 \dots$$

$$\left| \int \int_R f(x, y) dA - \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \right| \approx 1.5 \times 10^{-5}.$$

Example:  $\int \int_R \log(x + 2y) dA = 1.036 \dots$ ,  $n = 7, m = 5$

- ▶ Gaussian quadrature approximation

$$\int \int_R f(x, y) dA \approx \frac{(b-a)(d-c)}{4} \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \approx 1.03604817065 \dots$$

$$\left| \int \int_R f(x, y) dA - \frac{(b-a)(d-c)}{4} \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \right| \approx 6.4 \times 10^{-10}.$$

- ▶ Simpson rule approximation

$$\int \int_R f(x, y) dA \approx \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \approx 1.03603270963 \dots$$

$$\left| \int \int_R f(x, y) dA - \sum_{i=1}^n \sum_{j=1}^m c_i \hat{c}_j f(x_i, y_j) \right| \approx 1.5 \times 10^{-5}.$$

Gaussian quadrature much more accurate.

# Gaussian Quadrature: how good is it?

- ▶ Gaussian quadrature

$$\begin{aligned}\int_{-1}^1 f(x) dx &= c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n) \\ &\quad + O\left(\frac{4^{-n} |f^{(2n)}(\xi)|}{(2n)!}\right) \\ &\approx c_1 f(x_1) + c_2 f(x_2) + \cdots + c_n f(x_n),\end{aligned}$$

where  $x_1, x_2, \dots, x_n \in (-1, 1)$  are distinct roots of Legendre Polynomial  $P_n(x)$ .

- ▶ Error tiny for large  $n$  and  $|f^{(2n)}(\xi)| = O(1)$ .

**Improper Integral:** what if  $\max_{x \in [-1,1]} |f^{(2n)}(x)| \gg 1$ ?


Gaussian Quadrature for  $\int_{-1}^1 f(x)dx$ ,  $f(x) = \frac{1}{\sqrt{1-x^2}}$

$n$	Quadrature Value
10	2.9758
50	3.1071
250	3.1346
1000	3.1399

Gaussian Quadrature for  $\int_{-1}^1 f(x)dx$ ,  $f(x) = \frac{1}{\sqrt{1-x^2}}$

$n$	Quadrature Value
10	2.9758
50	3.1071
250	3.1346
1000	3.1399


- $f(x) = \frac{1}{\sqrt{1-x^2}}$  not smooth in  $[-1, 1]$



Gaussian Quadrature for  $\int_{-1}^1 f(x)dx$ ,  $f(x) = \frac{1}{\sqrt{1-x^2}}$

$n$	Quadrature Value
10	2.9758
50	3.1071
250	3.1346
1000	3.1399

- $f(x) = \frac{1}{\sqrt{1-x^2}}$  not smooth in  $[-1, 1]$



$$\int_{-1}^1 \frac{dx}{\sqrt{1-x^2}} \stackrel{x=\sin\theta}{=} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} 1 d\theta = 3.1416\dots$$

Improper Integral:  $\int_{-1}^1 \frac{f(x)}{\sqrt{1-x^2}} dx$

- ▶ Change of variable:

$$x = \sin \theta, \quad \theta \in [-\frac{\pi}{2}, \frac{\pi}{2}].$$

- ▶

$$dx = \cos \theta, \quad \sqrt{1-x^2} = \sqrt{1-\sin^2 \theta} = \cos \theta.$$

- ▶ New integral becomes proper.

$$\int_{-1}^1 \frac{f(x)}{\sqrt{1-x^2}} dx = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} f(\sin \theta) d\theta.$$

Improper Integral:  $\int_{-1}^1 \frac{f(x)}{\sqrt{1-x^2}} dx$

- ▶ Change of variable:

$$x = \sin \theta, \quad \theta \in [-\frac{\pi}{2}, \frac{\pi}{2}].$$

- ▶  
 $dx = \cos \theta, \quad \sqrt{1-x^2} = \sqrt{1-\sin^2 \theta} = \cos \theta.$
- ▶ New integral becomes proper.


$$\int_{-1}^1 \frac{f(x)}{\sqrt{1-x^2}} dx = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} f(\sin \theta) d\theta.$$

- ▶ **Example:** for  $f(x) = x^2$ , with 20-point Gaussian quadrature:

$$\int_{-1}^1 \frac{x^2}{\sqrt{1-x^2}} dx = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 \theta d\theta \approx 1.57079632679490,$$

accurate to 15 digits.


Improper Integral:  $\int_a^b f(x)dx$ ,  $f(x) = \frac{g(x)}{(x-a)^p}$ , for  $p < 1$



- Integral is improper if  $g(a) \neq 0$ :

$$\lim_{x \rightarrow a} \left| \frac{g(x)}{(x - a)^p} \right| = \infty.$$

Improper Integral:  $\int_a^b f(x)dx$ ,  $f(x) = \frac{g(x)}{(x-a)^p}$ , for  $p < 1$



- Integral is improper if  $g(a) \neq 0$ :

$$\lim_{x \rightarrow a} \left| \frac{g(x)}{(x-a)^p} \right| = \infty.$$

- Integral is not defined for  $p \geq 1$ :

$$\begin{aligned}\int_a^b \frac{1}{(x-a)^p} dx &= \lim_{M \rightarrow a^+} \int_M^b \frac{1}{(x-a)^p} dx \\ &= \lim_{M \rightarrow a^+} \frac{(x-a)^{1-p}}{1-p} \Big|_M^b = \frac{(b-a)^{1-p}}{1-p}.\end{aligned}$$

## Proper Method for improper Integral $\int_a^b \frac{g(x)}{(x-a)^p} dx$

- ▶ Assume a Taylor expansion on  $g(x)$ :

$$g(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(n)}(a)}{n!}(x-a)^n + \cdots .$$

## Proper Method for improper Integral $\int_a^b \frac{g(x)}{(x-a)^p} dx$

- ▶ Assume a Taylor expansion on  $g(x)$ :

$$g(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(n)}(a)}{n!}(x-a)^n + \cdots .$$

- ▶ Choose a  $(k+1)$ -term approximation:

$$P_k(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(k)}(a)}{k!}(x-a)^k.$$

## Proper Method for improper Integral $\int_a^b \frac{g(x)}{(x-a)^p} dx$

- ▶ Assume a Taylor expansion on  $g(x)$ :

$$g(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(n)}(a)}{n!}(x-a)^n + \cdots.$$

- ▶ Choose a  $(k+1)$ -term approximation:

$$P_k(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(k)}(a)}{k!}(x-a)^k.$$

- ▶ Improper integral decomposition

$$\int_a^b \frac{g(x)}{(x-a)^p} dx = \int_a^b \frac{P_k(x)}{(x-a)^p} dx + \int_a^b \frac{g(x) - P_k(x)}{(x-a)^p} dx.$$

## Proper Method for improper Integral $\int_a^b \frac{g(x)}{(x-a)^p} dx$

- ▶ Assume a Taylor expansion on  $g(x)$ :

$$g(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(n)}(a)}{n!}(x-a)^n + \cdots.$$

- ▶ Choose a  $(k+1)$ -term approximation:

$$P_k(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(k)}(a)}{k!}(x-a)^k.$$

- ▶ Improper integral decomposition

$$\int_a^b \frac{g(x)}{(x-a)^p} dx = \int_a^b \frac{P_k(x)}{(x-a)^p} dx + \int_a^b \frac{g(x) - P_k(x)}{(x-a)^p} dx.$$

- ▶ First integral is a simple sum:

$$\int_a^b \frac{P_k(x)}{(x-a)^p} dx = \sum_{j=0}^k \int_a^b \frac{g^{(j)}(a)}{j!} \frac{(x-a)^j}{(x-a)^p} dx = \sum_{j=0}^k \frac{g^{(j)}(a)}{j!(j+1-p)} (b-a)^{j+1-p}$$

## Proper Method for improper Integral $\int_a^b \frac{g(x)}{(x-a)^p} dx$

- ▶ Assume a Taylor expansion on  $g(x)$ :

$$g(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(n)}(a)}{n!}(x-a)^n + \cdots.$$

- ▶ Choose a  $(k+1)$ -term approximation:

$$P_k(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(k)}(a)}{k!}(x-a)^k.$$

- ▶ Improper integral decomposition

$$\int_a^b \frac{g(x)}{(x-a)^p} dx = \int_a^b \frac{P_k(x)}{(x-a)^p} dx + \int_a^b \frac{g(x) - P_k(x)}{(x-a)^p} dx.$$

- ▶ Second integral is a proper integral for  $k \gg p$ :

$$\int_a^b \frac{g(x) - P_k(x)}{(x-a)^p} dx = \int_a^b (x-a)^{k+1-p} \left( \sum_{j=k+1}^{\infty} \frac{g^{(j)}(a)}{j!} (x-a)^{j-k-1} \right) dx.$$

# Summary of Proper Method

- ▶ Choose a  $(k + 1)$ -term approximation:

$$P_k(x) = g(a) + g'(a)(x-a) + \frac{g''(a)}{2}(x-a)^2 + \cdots + \frac{g^{(k)}(a)}{k!}(x-a)^k.$$

- ▶ Improper integral decomposition

$$\begin{aligned} \int_a^b \frac{g(x)}{(x-a)^p} dx &= \int_a^b \frac{P_k(x)}{(x-a)^p} dx + \int_a^b \frac{g(x) - P_k(x)}{(x-a)^p} dx. \\ &= \left( \sum_{j=0}^k \frac{g^{(j)}(a)}{j!(j+1-p)} (b-a)^{j+1-p} \right) + \int_a^b G(x) dx, \end{aligned}$$

where  $G(x) = \begin{cases} 0, & \text{if } x = a, \\ \frac{g(x)-P_k(x)}{(x-a)^p}, & \text{if } x > a. \end{cases}$

Only require  $g(a), g'(a), \dots, g^{(k)}(a)$

Example:  $\int_0^1 \frac{e^x}{\sqrt{x}} dx$

- Take a 5-term Taylor expansion

$$P_4(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24}.$$

- 

$$\begin{aligned}\int_0^1 \frac{P_4(x)}{\sqrt{x}} dx &= \int_0^1 \left( \frac{1}{\sqrt{x}} + \sqrt{x} + \frac{x^{3/2}}{2} + \frac{x^{5/2}}{6} + \frac{x^{7/2}}{24} \right) dx \\ &= 2 + \frac{2}{3} + \frac{1}{5} + \frac{1}{21} + \frac{1}{108} \\ &\approx 2.9235450.\end{aligned}$$

- Composite Simpson's rule with  $n = 4, h = \frac{1}{n} = 0.25$  on

$$\int_0^1 G(x) dx, \quad \text{where } G(x) = \begin{cases} 0, & \text{if } x = 0, \\ \frac{e^x - P_4(x)}{\sqrt{x}}, & \text{if } x > 0. \end{cases}$$

Example:  $\int_0^1 \frac{e^x}{\sqrt{x}} dx$

$x$	$G(x)$
0.0	0
0.25	0.0000170
0.50	0.0004013
0.75	0.0026026
1.0	0.0099485

- Composite Simpson rule, with  $n = 4, h = \frac{1}{n} = 0.25$

$$\begin{aligned}\int_0^1 G(x)dx &\approx \frac{0.25}{3} (0 + 4 \times 0.0000170 + 2 \times 0.0004013 \\ &\quad + 4 \times 0.0026026 + 0.0099485) \approx 0.0017691.\end{aligned}$$

Example:  $\int_0^1 \frac{e^x}{\sqrt{x}} dx$

$x$	$G(x)$
0.0	0
0.25	0.0000170
0.50	0.0004013
0.75	0.0026026
1.0	0.0099485

- Composite Simpson rule, with  $n = 4, h = \frac{1}{n} = 0.25$

$$\begin{aligned}\int_0^1 G(x) dx &\approx \frac{0.25}{3} (0 + 4 \times 0.0000170 + 2 \times 0.0004013 \\ &\quad + 4 \times 0.0026026 + 0.0099485) \approx 0.0017691.\end{aligned}$$


- Improper integral decomposition

$$\begin{aligned}\int_0^1 \frac{e^x}{\sqrt{x}} dx &= \int_0^1 \frac{P_4(x)}{\sqrt{x}} dx + \int_0^1 \frac{e^x - P_4(x)}{\sqrt{x}} dx \\ &\approx 2.9235450 + 0.0017691 = 2.9253141.\end{aligned}$$

Improper Integral:  $\int_a^b f(x)dx$ ,  $f(x) = \frac{g(x)}{(b-x)^p}$ , for  $p < 1$

- ▶ Change of variable:  $z = -x$
- ▶ Left endpoint improper Integral

$$\int_a^b \frac{g(x)}{(b-x)^p} dx = \int_{-b}^{-a} \frac{g(-z)}{(z-(-b))^p} dz$$



Improper Integral:  $\int_a^\infty f(x)dx$ ,  $f(x) = \frac{g(x)}{x^p}$ , for  $p > 1$

- Integral is not defined for  $p \leq 1$ :

$$\begin{aligned}\int_a^\infty \frac{1}{x^p} dx &= \lim_{M \rightarrow \infty} \int_a^M \frac{1}{x^p} dx \\ &= \lim_{M \rightarrow \infty} \frac{x^{1-p}}{1-p} \Big|_a^\infty = \frac{a^{1-p}}{p-1}.\end{aligned}$$

- In general, change of variable  $z = \frac{1}{x}$ , assuming  $a > 0$ .

$$dx = -\frac{dz}{z^2}.$$

- Left end improper integral:

$$\int_a^\infty \frac{g(x)}{x^p} dx = \int_0^{\frac{1}{a}} g\left(\frac{1}{z}\right) z^{p-2} dz.$$

Example:  $\int_1^\infty f(x)dx$ ,  $f(x) = \frac{\sin(\frac{1}{x})}{x^{3/2}}$

- ▶ Change of variable  $z = \frac{1}{x}$ :

$$\int_1^\infty \frac{\sin(\frac{1}{x})}{x^{3/2}} dx = \int_0^1 \sin(z) z^{3/2} z^{-2} dz = \int_0^1 \frac{\sin(z)}{\sqrt{z}} dz.$$

Example:  $\int_1^\infty f(x)dx$ ,  $f(x) = \frac{\sin(\frac{1}{x})}{x^{3/2}}$

- ▶ Change of variable  $z = \frac{1}{x}$ :

$$\int_1^\infty \frac{\sin(\frac{1}{x})}{x^{3/2}} dx = \int_0^1 \sin(z) z^{3/2} z^{-2} dz = \int_0^1 \frac{\sin(z)}{\sqrt{z}} dz.$$

- ▶ Choose 5-term Taylor expansion on  $\sin(z)$ :  $P_4(z) = z - \frac{z^3}{6}$ .

Example:  $\int_1^\infty f(x)dx$ ,  $f(x) = \frac{\sin(\frac{1}{x})}{x^{3/2}}$

- ▶ Change of variable  $z = \frac{1}{x}$ :

$$\int_1^\infty \frac{\sin(\frac{1}{x})}{x^{3/2}} dx = \int_0^1 \sin(z) z^{3/2} z^{-2} dz = \int_0^1 \frac{\sin(z)}{\sqrt{z}} dz.$$

- ▶ Choose 5-term Taylor expansion on  $\sin(z)$ :  $P_4(z) = z - \frac{z^3}{6}$ .
- ▶

$$\begin{aligned}\int_0^1 \frac{\sin(z)}{\sqrt{z}} dz &= \int_0^1 \frac{z - \frac{z^3}{6}}{\sqrt{z}} dz + \int_0^1 \frac{\sin(z) - \left(z - \frac{z^3}{6}\right)}{\sqrt{z}} dz \\ &\approx 0.61904761 + \int_0^1 \frac{\sin(z) - \left(z - \frac{z^3}{6}\right)}{\sqrt{z}} dz.\end{aligned}$$

- ▶ Composite Simpson's rule with  $n = 16$ :

$$\int_0^1 \frac{\sin(z) - \left(z - \frac{z^3}{6}\right)}{\sqrt{z}} dz \approx 0.0014890097.$$

Example:  $\int_1^\infty f(x)dx$ ,  $f(x) = \frac{\sin(\frac{1}{x})}{x^{3/2}}$




$$\begin{aligned}\int_0^1 \frac{\sin(z)}{\sqrt{z}} dz &= \int_0^1 \frac{z - \frac{z^3}{6}}{\sqrt{z}} dz + \int_0^1 \frac{\sin(z) - \left(z - \frac{z^3}{6}\right)}{\sqrt{z}} dz \\ &\approx 0.61904761 + \int_0^1 \frac{\sin(z) - \left(z - \frac{z^3}{6}\right)}{\sqrt{z}} dz \\ &\approx 0.61904761 + 0.0014890097 \\ &= 0.62053661,\end{aligned}$$

accurate to 8-digit.

# Initial Value ODE

- ▶ The motion of a swinging pendulum

$$\frac{d^2\theta}{dt^2} + \frac{g}{L} \sin \theta = 0,$$




- ▶ Initial Value conditions

$$\theta(t_0) = \theta_0, \quad \text{and} \quad \theta'(t_0) = \theta'_0.$$

# Initial Value ODE

- ▶ The motion of a swinging pendulum

$$\frac{d^2\theta}{dt^2} + \frac{g}{L} \sin \theta = 0,$$



- ▶ Initial Value conditions


$$\theta(t_0) = \theta_0, \quad \text{and} \quad \theta'(t_0) = \theta'_0.$$

When does ODE have a solution?

# Initial Value ODE

- ▶ The motion of a swinging pendulum

$$\frac{d^2\theta}{dt^2} + \frac{g}{L} \sin \theta = 0,$$



- ▶ Initial Value conditions

$$\theta(t_0) = \theta_0, \quad \text{and} \quad \theta'(t_0) = \theta'_0.$$

When does ODE have a solution? How to compute it?

## Lipschitz condition

**Definition:** function  $f(t, y)$  satisfies a **Lipschitz condition** in the variable  $y$  on a set  $D \subset \mathbf{R}^2$  if a constant  $L > 0$  exists with

$$|f(t, y_1) - f(t, y_2)| \leq L |y_1 - y_2|,$$

whenever  $(t, y_1), (t, y_2)$  are in  $D$ .  $L$  is Lipschitz constant.

- ▶ **Example 1:** Show that  $f(t, y) = t|y|$  satisfies a Lipschitz condition on the region

$$D = \{(t, y) \mid 0 \leq t \leq T\}.$$

**Solution:** For any  $(t, y_1), (t, y_2)$  in  $D$ ,

$$|f(t, y_1) - f(t, y_2)| = |t|y_1| - t|y_2|| \leq t |y_1 - y_2| \leq L |y_1 - y_2|,$$

for  $L = T$ .

## Lipschitz condition

**Definition:** function  $f(t, y)$  satisfies a **Lipschitz condition** in the variable  $y$  on a set  $D \subset \mathbf{R}^2$  if a constant  $L > 0$  exists with

$$|f(t, y_1) - f(t, y_2)| \leq L |y_1 - y_2|,$$

whenever  $(t, y_1), (t, y_2)$  are in  $D$ .  $L$  is Lipschitz constant.

- ▶ **Example 2:** Show that  $f(t, y) = t y^2$  does not satisfy any Lipschitz condition on the region

$$D = \{(t, y) \mid 0 \leq t \leq T\}.$$

**Solution:** Choose  $(T, y_1), (T, y_2)$  in  $D$  with  $y_1 = 0, y_2 > 0$ ,

$$\frac{|f(T, y_1) - f(T, y_2)|}{|y_1 - y_2|} = T y_2,$$


which can be larger than  $L$  for any fixed  $L > 0$ .

## Convex Set

**Definition:** A set  $D \subset \mathbf{R}^2$  is convex if

whenever  $(t_1, y_1)$  and  $(t_2, y_2) \in D$

→ line segment  $(1 - \lambda)(t_1, y_1) + \lambda(t_2, y_2) \in D$  for all  $\lambda \in [0, 1]$ .



**Theorem:** Suppose  $f(t, y)$  is defined on a convex set  $D \subset \mathbf{R}^2$ . If a constant  $L > 0$  exists with

$$\left| \frac{\partial f}{\partial y}(t, y) \right| \leq L, \quad \text{for all } (t, y) \in D,$$

then  $f$  satisfies a Lipschitz condition with Lipschitz constant  $L$ .

**Theorem:** Suppose  $f(t, y)$  is defined on a convex set  $D \subset \mathbf{R}^2$ . If a constant  $L > 0$  exists with

$$\left| \frac{\partial f}{\partial y}(t, y) \right| \leq L, \quad \text{for all } (t, y) \in D,$$

then  $f$  satisfies a Lipschitz condition with Lipschitz constant  $L$ .

- ▶ **Example 1:** Show that  $f(t, y) = t y^2$  satisfies Lipschitz condition on the region

$$D = \{(t, y) \mid 0 \leq t \leq T, -Y \leq y \leq Y\}.$$

**Solution:**

$$\frac{\partial f}{\partial y}(t, y) = 2ty, \quad \left| \frac{\partial f}{\partial y}(t, y) \right| \leq 2T|y| \quad \text{for all } (t, y) \in D.$$

so  $f(t, y) = t y^2$  satisfies Lipschitz condition with  $L = 2T|Y|$ .

## What is going on with $f(t, y) = t y^2$ ?

- ▶  $f(t, y) = t y^2$  satisfies Lipschitz condition on the region

$$D = \{(t, y) \mid 0 \leq t \leq T, -Y \leq y \leq Y\}.$$

- ▶  $f(t, y) = t y^2$  doesn't satisfy Lipschitz condition on region

$$D = \{(t, y) \mid 0 \leq t \leq T\}.$$

Initial value problem

$$y'(t) = t y^2(t), \quad y(t_0) = \alpha > 0$$

has unique, but unbounded solution

$$y(t) = \frac{2\alpha}{2 + \alpha(t_0^2 - t^2)},$$

the denominator of which vanishes at

$$t = \sqrt{\frac{2}{\alpha} + t_0^2}.$$

# What is going on with $f(t, y) = t y^2$ ?

Initial value problem

$$y'(t) = t y^2(t), \quad y(t_0) = \alpha > 0$$

has unique, but unbounded solution

$$y(t) = \frac{2\alpha}{2 + \alpha(t_0^2 - t^2)},$$

the denominator of which vanishes at

$$t = \sqrt{\frac{2}{\alpha} + t_0^2}.$$

- ▶ for  $|t_0| < T$ , ODE has unique solution on

$$D = \{(t, y) \mid 0 \leq t \leq T, -Y \leq y \leq Y\}.$$

- ▶ for  $\sqrt{\frac{2}{\alpha} + t_0^2} < T$  ODE solution breaks down at  $t = \sqrt{\frac{2}{\alpha} + t_0^2}$  on

$$D = \{(t, y) \mid 0 \leq t \leq T\}.$$