

A SPLITTING THEOREM FOR n -*REA* DEGREES

RICHARD A. SHORE AND THEODORE A. SLAMAN

ABSTRACT. We prove that, for any D , A and U with $D >_T A \oplus U$ and r.e. in $A \oplus U$, there are pairs X_0, X_1 and Y_0, Y_1 such that $D \equiv_T X_0 \oplus X_1$; $D \equiv_T Y_0 \oplus Y_1$; and, for any i and j from $\{0, 1\}$ and any set B , if $X_i \oplus A \geq_T B$ and $Y_j \oplus A \geq_T B$ then $A \geq_T B$. We then deduce that for any degrees \mathbf{d} , \mathbf{a} , and \mathbf{b} such that \mathbf{a} and \mathbf{b} are recursive in \mathbf{d} , $\mathbf{a} \not\leq_T \mathbf{b}$, and \mathbf{d} is n -*REA* in \mathbf{a} , \mathbf{d} can be split over \mathbf{a} avoiding \mathbf{b} . This shows that the Main Theorem of Cooper [1990] and [1993] is false.

1. INTRODUCTION

The basic formal definitions of computable sets and functions were proposed independently, and in various equivalent forms, in the 30s by many researchers. The standard model now is that of the Turing machine. Turing [1939] also introduced the notion of relative computability to capture the idea of one set A being easier to compute than another B in the sense that if we knew how to settle membership questions in B we could do the same for A . The model here consists of Turing machines with oracles, i.e. black boxes that supply the required answers about B . We say that A is computable from (or recursive in) B if there is a Turing machine which, when equipped with an oracle for B , computes (the characteristic function of) A , i.e. for some e , $\phi_e^B = A$. (Here we are assuming a standard listing ϕ_e^B of the Turing machines with oracles for B .) We denote this relation by $A \leq_T B$ which we read as A is (Turing) reducible to B or A is recursive (computable) in B . This relation is transitive and reflexive and so induces an equivalence relation \equiv_T ($A \equiv_T B \Leftrightarrow A \leq_T B \wedge B \leq_T A$) and a partial order also denoted by \leq_T on the equivalence classes. These equivalence classes are called (Turing) degrees and the equivalence class of a set $A \subseteq \omega$ is called its degree. It is typically denoted by \mathbf{a} or $\text{deg}(A)$.

The structure \mathcal{D} of these degrees has been the object of extensive study over the past fifty or sixty years. (A survey from the early 80s is Shore [1985]. Current ones can be found in Griffor [1999].) At first, attention focused on basic structural questions about \mathcal{D} such as determining which partial orders or lattices can be embedded in \mathcal{D} and characterizing its initial segments. For the past twenty years a considerable amount of attention has been paid to more global issues such as automorphisms of, and definability in, \mathcal{D} . Of course, some of the vast store of structural information discovered earlier on played important roles in these later investigations. Particularly important examples in the early work on global issues

1991 *Mathematics Subject Classification*. Primary 03D25, 03D39; Secondary 03D55, 03E45.

Key words and phrases. degrees, Turing degrees, recursively enumerable degrees, splitting theorems.

Partially supported by NSF Grant DMS-9802843.

Partially supported by NSF Grant DMS-97-96121.

include some of the results on initial segments and Spector's exact pair theorem that says that every countable ideal is the intersection of two principle ideals. More specific local structural results that have played central roles in definability results for \mathcal{D} include ones revolving around density, or the lack thereof, and the interactions between certain special subclasses of degrees related to recursive enumerability and all of the degrees. (A degree \mathbf{b} is *recursively enumerable relative to* (or *in*) a degree \mathbf{a} if there is a set $B \in \mathbf{b}$ which can be enumerated by a function recursive in an $A \in \mathbf{a}$, i.e. B is the range of some ϕ_e^A .)

In particular, Jockusch and Shore [1984] used two such facts due to Sacks [1963] and an analysis of operators corresponding to iterations of recursive enumerability and ones corresponding to sets which are effective Boolean combinations of r.e. ones to show that the class \mathcal{A} of degrees of arithmetic sets (those definable in arithmetic) is definable in \mathcal{D} . The basic structural facts needed were that no r.e. degree \mathbf{a} can be a minimal cover of any degree \mathbf{b} , i.e. if $\mathbf{b} < \mathbf{a}$ and \mathbf{a} is r.e. then there is always a degree \mathbf{c} strictly between \mathbf{b} and \mathbf{a} and that every degree \mathbf{b} has a minimal cover \mathbf{a} which is the degree of a set which is an effective Boolean combination of sets r.e. in B of a particular form.

In a similar vein, Cooper proposed an approach to defining $\mathbf{0}'$, the degree of the halting problem, and so the jump operator taking \mathbf{a} to \mathbf{a}' . (Here \mathbf{a}' is the degree of the halting problem for machines with an oracle for A , i.e. $\mathbf{a}' = \text{deg}(A')$ where $A' = \{e \mid \text{the computation of } \phi_e^A(e) \text{ eventually halts}\}$.) The structural property he considered was considerably more complicated than the nonminimality one of r.e. degrees used by Jockusch and Shore but was also based, in one direction, on an old result of Sacks'. Here is Cooper's property, some associated classes of degrees and the relevant theorem of Sacks corresponding to the nonminimality results for r.e. degrees.

Definition 1.1. \mathbf{d} is splittable over \mathbf{a} avoiding \mathbf{b} if either $\mathbf{a}, \mathbf{b} \not\leq \mathbf{d}$ or $\mathbf{b} \leq \mathbf{a}$ or there are $\mathbf{d}_0, \mathbf{d}_1$ such that $\mathbf{a} <_T \mathbf{d}_0, \mathbf{d}_1 <_T \mathbf{d}$, $\mathbf{d}_0 \vee \mathbf{d}_1 = \mathbf{d}$ and $\mathbf{b} \not\leq_T \mathbf{d}_0, \mathbf{d}_1$. $\mathcal{C}_1 = \{\mathbf{c} \mid \forall \mathbf{a}, \mathbf{b} (\mathbf{a} \vee \mathbf{c} \text{ is splittable over } \mathbf{a} \text{ avoiding } \mathbf{b})\}$. $\overline{\mathcal{C}}_1 = \{\mathbf{d} \mid \exists \mathbf{c} \in \mathcal{C}_1 (\mathbf{d} \leq_T \mathbf{c})\}$.

Theorem 1.2. (Sacks [1963]) *Every r.e. degree \mathbf{d} is in \mathcal{C}_1 .*

The analog of the Sacks' minimal cover theorem in Cooper's work was his Main Theorem that there is a set C which is $d - r.e.$, i.e. the set-difference of two r.e. sets, (and so a corresponding operator) such that there are \mathbf{a} and \mathbf{b} such that $\mathbf{c} \equiv_T \text{deg}(C)$ is not splittable over \mathbf{a} avoiding \mathbf{b} . Together with a general theorem on the range of such operators and their interaction under join with degrees not below $\mathbf{0}'$, this result would have sufficed to define $\mathbf{0}'$ and so, by relativization, the jump operator itself. Cooper [1990] proposed to argue for the existence of such a set with a $0'''$ priority construction similar to that proving the Nonsplitting Theorem of Lachlan [1975] for the recursively enumerable degrees. In this paper, we prove a splitting theorem (Theorem 2.3) a special case of which shows that there is no such set C and so Cooper's proposed property does not define the jump. In Shore and Slaman [2000] we provide a very different definition of $\mathbf{0}'$ and the jump operator.

2. THE SPLITTING THEOREM

We begin with the definition of the iteration of recursive enumerability basic to the work of both Jockusch and Shore and Cooper.

Definition 2.1. A set W is 1 -*REA* relative to, or in, A if it is *r.e.* in A and above it in Turing degree. W is $(n+1)$ -*REA* in A if it is 1 -*REA* relative to some U which is n -*REA* in A . A degree \mathbf{w} is n -*REA* relative to, or in, \mathbf{a} if it contains a set W which is n -*REA* relative to some set $A \in \mathbf{a}$. If $\mathbf{a} = \mathbf{0}$, we just say that W and \mathbf{w} are n -*REA*.

As Lachlan has shown that every d -*r.e.* set is of 2 -*REA* degree (see Jockusch and Shore [1984] for a proof of a more general fact), it suffices to prove that every 2 -*REA* degree \mathbf{d} is splittable over every \mathbf{a} avoiding any \mathbf{b} to show that there is no d -*r.e.* set as claimed in the Main Theorem of Cooper [1990], [1993]. In fact, we prove this for every n -*REA* degree \mathbf{d} . The proof is not uniform in the sense that given D , A and B we do not construct a splitting (D_0, D_1) of D that lies above A and not above B in such a way as to effectively produce indices for the D_i from those of the given sets. Instead, we produce various candidates some pair of which constitute the desired splitting. The case for $n = 1$ is the Sacks splitting theorem (Theorem 1.2). We begin with a result that suffices for the case $n = 2$ and then use it in an induction argument to get the full result.

Theorem 2.2. Let D , A and U be given so that $D >_T A \oplus U$ and D is recursively enumerable in $A \oplus U$. Then there are pairs X_0, X_1 and Y_0, Y_1 with the following properties.

- (1) $D \equiv_T X_0 \oplus X_1$ and $D \equiv_T Y_0 \oplus Y_1$.
- (2) For any $i, j \in \{0, 1\}$, and any set B , if $X_i \oplus A \geq_T B$ and $Y_j \oplus A \geq_T B$ then $A \geq_T B$.

The desired splitting theorem for n -*REA* degrees is an easy corollary of Theorem 2.2.

Theorem 2.3. Suppose that \mathbf{d} , \mathbf{a} , and \mathbf{b} are given and \mathbf{d} is n -*REA* in \mathbf{a} . Then \mathbf{d} can be split over \mathbf{a} avoiding \mathbf{b} .

Proof. Without loss of generality we may assume that \mathbf{a} and \mathbf{b} are recursive in \mathbf{d} and $\mathbf{a} \not\geq_T \mathbf{b}$. We deduce Theorem 2.3 from Theorem 2.2 by an induction on n . Let D , A , and B be representatives of \mathbf{d} , \mathbf{a} , and \mathbf{b} , respectively.

When n is equal to 1, D is recursively enumerable in A . By the Sacks Splitting Theorem (Theorem 1.2) relativized to A , \mathbf{d} can be split over \mathbf{a} avoiding \mathbf{b} .

Now assume that n is greater than 1 and assume that Theorem 2.3 holds for all numbers less than n . Since \mathbf{d} is n -*REA* in \mathbf{a} , there is a set U such that U is $(n-1)$ -*REA* relative to A , U is recursive in D and D is recursively enumerable in U .

First, if D is recursive in U , then D has the same degree as U and the inductive hypothesis verifies the claim of Theorem 2.3. So, assume D is not recursive in U . Then $D >_T A \oplus U$ and is recursively enumerable in $A \oplus U$. Thus we can apply Theorem 2.2 to get X_0, X_1 and Y_0, Y_1 as indicated in its conclusion. If neither $A \oplus X_0$ nor $A \oplus X_1$ computes B , then the degrees of $A \oplus X_0$ and $A \oplus X_1$ form a splitting of \mathbf{d} over \mathbf{a} avoiding \mathbf{b} . Otherwise, B is recursive in either $A \oplus X_0$ or $A \oplus X_1$. By the second condition in Theorem 2.2, B cannot be recursive in either $A \oplus Y_0$ or $A \oplus Y_1$. Consequently, the degrees of $A \oplus Y_0$ and $A \oplus Y_1$ form a splitting of \mathbf{d} over \mathbf{a} avoiding \mathbf{b} . Thus, in either case, \mathbf{d} can be split over \mathbf{a} avoiding \mathbf{b} as required. \square

The remainder of this paper is devoted to proving Theorem 2.2. As in the statement of Theorem 2.2, let D , A and U be given so that $D >_T A \oplus U$ and D is recursively enumerable relative to $A \oplus U$. We construct pairs X_0, X_1 and Y_0, Y_1 to satisfy the conclusions of the theorem.

In the following, we will identify sets with their characteristic functions. For example, $D(n) = 0$ is synonymous with $n \notin D$, and $D(n) = 1$ is synonymous with $n \in D$. We first describe the requirements that we need to satisfy and then provide the construction of the desired sets from D and finally the verifications that the requirements are satisfied..

2.1. Join requirements. We directly ensure that $X_0 \oplus X_1$ and $Y_0 \oplus Y_1$ compute D by enforcing the following conditions.

$$(2.1) \quad (\forall n)[n \in D \text{ if and only if } X_0(n) = X_1(n)]$$

$$(2.2) \quad (\forall n)[n \in D \text{ if and only if } Y_0(n) = Y_1(n)].$$

Note that we can specify X_i or Y_j arbitrarily and still define X_{1-i} or Y_{1-j} , respectively, to satisfy Equations 2.1 and 2.2.

2.2. Infima requirements. We ensure that our sets satisfy the second claim in Theorem 2.2 by a priority construction. We consider requirements P of the following form, in which i and j belong to $\{0, 1\}$, Φ and Ψ are Turing functionals, and B is a free variable.

$$(2.3) \quad \text{If } \Phi(X_i \oplus A) = \Psi(Y_j \oplus A) = B, \text{ then } A \geq_T B.$$

Let $(P_i : i \in \mathbb{N})$ be a recursive enumeration of all requirements of the above form. To fix our notation, let P_k be the requirement

$$(2.4) \quad \text{If } \Phi_k(X_{i_k} \oplus A) = \Psi_k(Y_{j_k} \oplus A) = B_k, \text{ then } A \geq_T B_k.$$

We organize our construction so that for all P_k , either one of $\Phi_k(X_{i_k} \oplus A)$ or $\Psi_k(Y_{j_k} \oplus A)$ is not total, they are not equal, or their common value is recursive in A . During the construction, we search for an argument at which we can make $\Phi_k(X_{i_k} \oplus A)$ and $\Psi_k(Y_{j_k} \oplus A)$ unequal. If no such opportunity appears and both functions are total, then we conclude that their common value must be recursive in A .

2.3. Constructing the X_i 's and Y_j 's recursively in D .

Definition 2.4. (1) *A finite condition on one of the X_i 's or Y_j 's is a 0-1 valued function whose domain is a finite initial segment of \mathbb{N} . That is, a finite condition on a set is a specification of a finite initial segment of the values of that set. Two conditions are compatible if they agree on the domain they have in common.*

- (2) *A P_k -split over A consists of a natural number n , a pair of finite conditions p_0 and p_1 , and a pair of computations in Φ_k and Ψ_k of lengths less than the domains of p_0 and p_1 such that $\Phi_k(n, p_0 \oplus A)$ and $\Psi_k(n, p_1 \oplus A)$ are defined by these computations and have different values.*
- (3) *We say that $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ extend the conditions in the above P_k -split if X_{i_k} extends p_0 and Y_{j_k} extends p_1 .*
- (4) *We say that $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ are compatible with the conditions in the above P_k -split if X_{i_k} and p_0 are compatible and Y_{j_k} and p_1 are compatible.*

Note, if $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ extend the conditions in a P_k -split, then $\Phi_k(X_{i_k} \oplus A)$ and $\Psi_k(Y_{j_k} \oplus A)$ take different values at the number n mentioned in that split.

In the following, we assume we have a universal enumeration over all k of all P_k splits over A .

Definition 2.5. (1) Let $D[s]$ denote the set of numbers less than or equal to s which are enumerated in D relative to $A \oplus U$ by computations of length less than or equal to s .

(2) For each $x \in \mathbb{N}$, let s_x be the least stage s such that $D[s]$ and D are equal on all numbers less than or equal to x , that is $D[s] \upharpoonright x \oplus 1 = D \upharpoonright x \oplus 1$.

We now compute the X_i 's and Y_j 's from D by recursion as follows.

- (1) To compute these sets at argument x , first compute their restrictions to all numbers less than x . If x is 0, then these restrictions are null functions.
- (2) Compute s_x .
- (3) Say that k requires attention at x if k is less than or equal to x and the following conditions hold.
 - (a) $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ do not extend the conditions in any P_k -split enumerated at or before stage s_x (of our fixed enumeration for the P_k splits over A).
 - (b) There is a P_k -split enumerated at or before stage s_x such that $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ are compatible with the conditions in that split.

If there is no k which requires attention at x , then let $X_0(x)$ and $Y_0(x)$ both equal 1 and let $X_1(x)$ and $Y_1(x)$ both equal $D(x)$. Thus, $X_0(x) = X_1(x)$ if and only if $Y_0(x) = Y_1(x)$ if and only if $D(x) = 1$.

If there is a k which requires attention at x , then let p_0 and p_1 be the conditions mentioned in the P_k -split which appeared first in our universal enumeration of splits among those P_k -splits of highest priority, i.e. with the smallest k , which are compatible with $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$. Define $X_{i_k}(x)$ and $Y_{j_k}(x)$ to equal $p_0(x)$ and $p_1(x)$, respectively, where those are defined. This leaves at least one of each pair $\{X_0, X_1\}$ and $\{Y_0, Y_1\}$ undefined at x . Define the sets whose values at x have not yet been determined to ensure the equivalences $X_0(x) = X_1(x)$ if and only if $Y_0(x) = Y_1(x)$ if and only if $D(x) = 1$.

Here is a short summary of the algorithm. Use the enumeration of D relative to $A \oplus U$ to produce a finite set of splits for various P_k 's, and then define the X_i 's and Y_j 's at x compatibly with the split for the highest priority requirement which requires attention.

Remark 2.6. The function mapping $D \upharpoonright x$ to s_{x-1} , $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ is recursive in $A \oplus U$.

Proof. Suppose we are given $D \upharpoonright x$ and by induction have computed $X_0 \upharpoonright x-2$, $X_1 \upharpoonright x-2$, $Y_0 \upharpoonright x-2$, and $Y_1 \upharpoonright x-2$. We can run the enumeration of D relative to $A \oplus U$ until the stage s_{x-1} at which each element of $D \upharpoonright x$ has been enumerated. The universal enumeration of P_k splits over A is recursive in A , and so we can compute the set of splits enumerated at or before stage s_{x-1} using $A \oplus U$. The values of the X_i 's and Y_j 's at $x-1$ are effectively determined from this collection of

splits and the value of D at $x - 1$, data which is now seen to be available to $A \oplus U$ from $D \upharpoonright x$. \square

Definition 2.7. Let $X_0[s]$, $X_1[s]$, $Y_0[s]$, and $Y_1[s]$ be the sets computed from $D[s]$ in the way that we specified that D computes X_0 , X_1 , Y_0 , and Y_1 .

2.4. Satisfaction of the requirements.

Lemma 2.8. For each k , there are only finitely many x such that P_k requires attention at x .

Proof. We verify Lemma 2.8 by a Friedberg-style priority argument. By induction on k , choose x_0 so that for all x greater than x_0 no requirement of index less than k requires attention at x . Suppose that P_k requires attention at some x_1 greater than x_0 . Then the values of X_{i_k} and Y_{j_k} at x will be defined so that $X_{i_k} \upharpoonright x \oplus 1$ and $Y_{j_k} \upharpoonright x \oplus 1$ are compatible with the conditions in the first P_k -split for which this is possible. Since no requirement of lower index can require attention, the same P_k -split will be used during subsequent stages until reaching an x_2 such that the X_i 's and Y_j 's restrictions to x_2 extend that P_k -split. But then P_k cannot require attention for any argument greater than x_2 . \square

Lemma 2.9. Suppose that $\Phi_k(X_{i_k} \oplus A) = \Psi_k(Y_{j_k} \oplus A)$ and their common value B_k is not recursive in A . Then for all x there is an s greater than x with the following properties.

- (1) $s \geq s_x$.
- (2) There is a P_k -split which is enumerated at or before stage s such that $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ are compatible with the conditions in that split.

Proof. For a contradiction, fix x so that there is no s which satisfies the conclusions of Lemma 2.9.

Then every t greater than s_x satisfies the first property specified in the conclusion of Lemma 2.9, so there cannot be any P_k -split such that $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ are compatible with the conditions in that split.

Consider any conditions p_0 and p_0^* extending $X_{i_k} \upharpoonright x$. If there were an n such that $\Phi_k(n, p_0 \oplus A)$ and $\Phi_k(n, p_0^* \oplus A)$ are defined and give incompatible values, then one of them could be paired with the initial segment of Y_{j_k} used to compute $\Psi_k(n, Y_{j_k} \oplus A)$ to produce a P_k -split.

Now, conclude that B_k is recursive in A : Given n , find p_0 extending $X_{i_k} \upharpoonright x$ such that $\Phi_k(n, p_0 \oplus A)$ is defined. The value of $\Phi_k(n, p_0 \oplus A)$ must equal $B_k(n)$, since it must equal $\Phi_k(n, X_{i_k} \oplus A)$. Thus, $A \geq_T B_k$ for the desired contradiction. \square

Lemma 2.10. For all k , if $\Phi_k(X_{i_k} \oplus A) = \Psi_k(Y_{j_k} \oplus A)$, then their common value is recursive in A .

Proof. For a contradiction, suppose that $\Phi_k(X_{i_k} \oplus A) = \Psi_k(Y_{j_k} \oplus A)$ and that their common value is not recursive in A . We will show that D is recursive in $A \oplus U$.

By Lemma 2.8, fix t_0 so that for all k^* less than or equal to k and all x greater than t_0 , P_{k^*} does not require attention at x .

Take $D \upharpoonright t_0$ as a given finite amount of data. We compute the value of D at arguments $x > t_0$ using the already computed $D \upharpoonright x$ as follows.

- (1) Compute s_{x-1} , $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$. (See Remark 2.6 and text preceding it for the effectiveness of this step relative to $A \oplus U$.)

- (2) Compute the least t such that the following conditions are satisfied.
- (a) t is greater than the maximum of x and s_{x-1} .
 - (b) There is a P_k -split which is enumerated at or before stage t such that $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ are compatible with the conditions in that split.

We let t_x denote this t .

- (3) Return the value of $D[t_x]$ at x as the value of D at x .

By Lemma 2.9, there is a stage t and a P_k -split which is enumerated at or before stage t such that $X_0 \upharpoonright x$, $X_1 \upharpoonright x$, $Y_0 \upharpoonright x$, and $Y_1 \upharpoonright x$ are compatible with the conditions in that split. Thus, the above procedure will find some t_x and will return some value for D at x .

We chose t_0 so that no strategy with index less than or equal to k requires attention after stage t_0 . In particular, P_k cannot require attention at x . But then, the P_k -split which is enumerated before or during stage t_x can not have been enumerated before or during stage s_x . Consequently, s_x is less than t_x and $D(x) = D[t_x](x)$. Thus, the calculation relative to $A \oplus U$ correctly returns the value of D at x .

The conclusion that D is recursive in $A \oplus U$ is the desired contradiction, which proves Lemma 2.10 and so Theorem 2.2. \square

REFERENCES

- Cooper, S. B. [1990], The jump is definable in the structure of the degrees of unsolvability (research announcement), *Bull. Am. Math. Soc.* **23**, 151-158.
- Cooper, S. B. [1993], On a conjecture of Kleene and Post, Department of Pure Mathematics, Leeds University, 1993 Preprint Series No. 7.
- Griffor, E. (ed.) [1999], *Handbook of Computability Theory*, North-Holland, Amsterdam.
- Jockusch, C. G. Jr. and Shore, R. A. [1984], Pseudo-jump operators II: transfinite iterations, hierarchies and minimal covers, *J. Symb. Logic* **49**, 1205-1236.
- Sacks, G. E. [1963], On the degrees less than $\mathbf{0}'$, *Ann. of Math. (2)* **77**, 211-231.
- Shore, R. A. 1985, The structure of the degrees of unsolvability, in *Recursion theory*, Proc. Symp. Pure Math. **42**, A. Nerode and R. A. Shore eds., Amer. Math. Soc., Providence RI, 33-51.
- Shore, R. A. and Slaman, T. A. [2000], Defining the Turing jump, to appear.
- Turing, A. M. [1939], Systems of logic based on ordinals, *Proc. London Math. Soc. (3)* **45**, 161-228.

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA NY 14853

E-mail address: shore@math.cornell.edu

URL: http://math.cornell.edu/~shore/

UNIVERSITY OF CALIFORNIA, BERKELEY, BERKELEY, CA 94720-3840,

E-mail address: slaman@math.berkeley.edu

URL: http://www.math.berkeley.edu/~slaman/